

SULLIVAN MINIMAL MODELS OF CLASSIFYING SPACES FOR NON-FORMAL SPACES OF SMALL RANK

HIROKAZU NISHINOBU AND TOSHIHIRO YAMAGUCHI

ABSTRACT. We consider certain rational homotopical conditions of simply connected CW complex X such that the rational cohomology of the classifying space $Baut_1X$ for fibrations with two-stage fibre X is (not) free. First, we consider when is $Baut_1X$ a rational factor of $Baut_1(X \times S^n)$ for an odd-integer n and observe for a non-formal elliptic space X of rank 3. Second, we compute the Sullivan minimal models of $Baut_1X$ when X are certain non-formal pure spaces of rank 5.

1. INTRODUCTION

Let X be a simply connected finite CW complex and aut_1X the identity component of self-homotopy equivalences of X . The Dold-Lashof classifying space [2], $Baut_1X$, is the classifying space for orientable fibrations with fiber the homotopy type of X . In 1968, J. Milnor [15] showed that, when $X = S^n$, $H^*(Baut_1X; \mathbb{Q}) \cong \mathbb{Q}[v]$ where $|v| = 2n$ if n is even and $|v| = n + 1$ if n is odd. Let $M(X)$ be the Sullivan minimal model [24] of X and $DerM(X)$ the DGL (differential graded Lie algebra) of the negative derivations on $M(X)$ (see §2 below). In 1977, D. Sullivan [24] indicated that $DerM(X)$ determines the rational homotopy type of $Baut_1X$, that is, $DerM(X)$ is a Quillen's DGL-model [17]. We are interested in the rational homotopical properties of it. (See [11] as a recent research of this direction.) It seems to be very difficult to expect understanding them immediately from the Sullivan model itself without the direct computation (see Theorem 2.1 below) of derivations of $M(X)$ et al. (cf. [19], [6]). Indeed, it is complicated even when X is a product of spheres [21]. For example, when X is $S^3 \times S^5 (\simeq_{\mathbb{Q}} SU(3))$, $Baut_1X$ is not a rational H-space and it is not even formal (see Definition 1.8 below). Then the rational cohomology is infinitely generated as

$$H^*(Baut_1X; \mathbb{Q}) \cong \mathbb{Q}[v] \otimes \Lambda(w_0, w_1, w_2, \dots) / (\{vw_i\}_i, \{w_iw_j\}_{i < j})$$

with $|v| (= \deg(v)) = 4$ and $|w_i| = 3 + 6i$ even though $H^*(Baut_1S^3; \mathbb{Q})$ and $H^*(Baut_1S^5; \mathbb{Q})$ are given as $\mathbb{Q}[v]$ and $\mathbb{Q}[u]$ with $|u| = 6$, respectively. A simply connected CW complex X is said to be *elliptic* if the dimensions of rational cohomology and homotopy are finite. Furthermore, an elliptic space X is said to be an F_0 -space if

$$H^*(X; \mathbb{Q}) \cong \mathbb{Q}[x_1, \dots, x_n] / (f_1, \dots, f_n),$$

in which $|x_i|$ are even and $f_1, \dots, f_n$ forms a regular sequence in $\mathbb{Q}[x_1, \dots, x_n]$. Then the Sullivan model is given as $M(X) = (\Lambda(x_1, \dots, x_n, y_1, \dots, y_n), d)$ with $|x_i|$ even and $|y_i|$ odd with $dx_i = 0$ and $dy_i = f_i \in \mathbb{Q}[x_1, \dots, x_n]$. In 1977, S. Halperin [9]

2010MSC: 55P62, 55R15

Keywords: rational homotopy theory, Sullivan (minimal) model, classifying space for fibration, formal, coformal, elliptic space, F_0 -space, pure space, derivation

conjectured that the Serre spectral sequences of all fibration $X \rightarrow E \rightarrow B$ collapse for any F_0 -space X . The Halperin's conjecture for an F_0 -space X is equivalent to

$$H^*(Baut_1 X; \mathbb{Q}) \cong \mathbb{Q}[v_1, \dots, v_m]$$

for some even-degrees' elements $v_1, \dots, v_m$ [13]. It is true when X is a homogeneous space G/H where G and H be compact connected Lie groups with H is a subgroup of G and $\text{rank} G = \text{rank} H$ due to Shiga-Tezuka [20]. Of course, even if a space X is not an F_0 -space, $H^*(Baut_1 X; \mathbb{Q})$ may be a polynomial algebra. For example, when X is $S^3 \times S^3$, $H^*(Baut_1 X; \mathbb{Q}) \cong \mathbb{Q}[v, u]$ with $|v| = |u| = 4$. (See [11] and [26] for the realization problem [3, p.519] of a polynomial algebra.) It is natural to ask

Question 1.1. *When is $H^*(Baut_1 X; \mathbb{Q})$ a polynomial algebra or free ?*

The meaning of “ $H^*(Baut_1 X; \mathbb{Q})$ is free” is that there is a graded algebra isomorphism $H^*(Baut_1 X; \mathbb{Q}) \cong \Lambda(x_1, \dots, x_m) \otimes \mathbb{Q}[y_{m+1}, \dots, y_n]$ with $|x_i| (= |y_i|) = k_i$ for some $m \leq n$ (i.e., the differential of Sullivan minimal model is zero). Then the rational homotopy set of classifying maps from a space Y is given as

$$[Y, (Baut_1 X)_{\mathbb{Q}}] = [Y, \Pi_{i=1}^n K(\mathbb{Q}, k_i)] = \bigoplus_{i=1}^n H^{k_i}(Y; \mathbb{Q}),$$

where $K(\mathbb{Q}, k_i)$ is the Eilenberg-Mac Lane space. That is, the tuple of an n -elements of the cohomology of Y rationally determine a fibration with fibre X and base Y . Here $Z_{\mathbb{Q}}$ means the rationalization of a nilpotent space Z .

In this paper, first, as an important obstruction for Problem 1.1, we pay attention to that $H^*(Baut_1(X \times Y); \mathbb{Q})$ can not be formulated by $H^*(Baut_1 X; \mathbb{Q})$ and $H^*(Baut_1 Y; \mathbb{Q})$ since $Baut_1 X$ is not even a rational factor of $Baut_1(X \times Y)$, i.e., $(Baut_1(X \times Y))_{\mathbb{Q}} \not\cong (Baut_1 X)_{\mathbb{Q}} \times C$ for any rational space C , in general. Refer S.B.Smith's works [21] and [22] in 2001. Thus our motivation is the following

Lemma 1.2. *If $H^*(Baut_1(X \times Y); \mathbb{Q})$ is free, then $H^*(Baut_1 X; \mathbb{Q})$ and $H^*(Baut_1 Y; \mathbb{Q})$ are both free. But the converse is not true in general.*

We prepare the following general notation to start our observation.

Definition 1.3. *A space S with a map $f : S \rightarrow T$ is said to have a rational retraction for T when there is a map $r : T_{\mathbb{Q}} \rightarrow S_{\mathbb{Q}}$ with $r \circ f_{\mathbb{Q}} \simeq id_{S_{\mathbb{Q}}}$. Also S is said to have a weak (rational) retraction for T when there is a graded Lie algebra map $R : \pi_*(\Omega T)_{\mathbb{Q}} \rightarrow \pi_*(\Omega S)_{\mathbb{Q}}$ with $R \circ \pi_*(\Omega f)_{\mathbb{Q}} = id_{\pi_*(\Omega S)_{\mathbb{Q}}}$.*

Thus we have “rational factor $\Rightarrow$ rational retraction $\Rightarrow$ weak retraction”. In particular, when a map $f : S \rightarrow T$ is $\pi_{\mathbb{Q}}$ -injective, “ $H^*(T; \mathbb{Q})$ is free $\Rightarrow S$ is a rational factor of T ” from Lemma 2.2 below. In general, a map $f : X \rightarrow Y$ does not induce a map between identity components of spaces of self-equivalences $aut_1 X \rightarrow aut_1 Y$. Though the inclusion $i_X : X \rightarrow X \times Y$ induces the map $j_X : aut_1 X \rightarrow aut_1(X \times Y)$ with $j_X(g) = g \times id_Y$, where id_Y is the identity of Y . Thus there is the morphism $Bj_X : Baut_1 X \rightarrow Baut_1(X \times Y)$. For example, $Baut_1 S^3$ and $Baut_1 S^5$ are not rational factors of $Baut_1(S^3 \times S^5)$. $Baut_1 S^3$ has a rational retraction for $Baut_1(S^3 \times S^5)$ but $Baut_1 S^5$ does not even a weak retraction. See §2 for the detail. The following question is a key to approach Question 1.1.

Question 1.4. *When is $Baut_1 X$ (or $Baut_1 Y$) a rational factor of $Baut_1(X \times Y)$? More generally, when does $Baut_1 X$ (or $Baut_1 Y$) have a rational (weak) retraction for $Baut_1(X \times Y)$?*

For example, recall S.B.Smith's

Theorem 1.5. ([21, Theorem 3]) *Suppose that X is an F_0 -space with $\text{Der}_+ H^*(X; \mathbb{Q}) = 0$ and Y is a rational H -space. If $\min \pi_*(X)_{\mathbb{Q}} + \min \pi_*(Y)_{\mathbb{Q}} \geq \max \pi_*(X \times Y)_{\mathbb{Q}}$ and $\max \pi_*(X)_{\mathbb{Q}} \leq \min \pi_*(Y)_{\mathbb{Q}}$, then*

$$\text{Baut}_1(X \times Y) \simeq_{\mathbb{Q}} \text{Baut}_1 X \times \|\text{Der}(H^*(Y; \mathbb{Q}), H^*(X \times Y; \mathbb{Q}))\|,$$

where $\|L\|$ is the spatial realization of a DGL L [17].

Futhermore there are some examples in [21, section 5]. In this paper, we consider a sufficient condition for having a weak retraction in the case that Y is an odd-sphere.

Definition 1.6. (a) *The (m, k) -evaluation of X :*

$$\mu_{m,k}(X) : H_m(\text{Der} M(X)) \otimes H^k(X; \mathbb{Q}) \rightarrow H_{m-k}(\text{Der} M(X))$$

is given by $\mu_{m,k}([\sigma] \otimes [w]) = [\sigma \otimes w]$. Here $([\sigma] \otimes [w])(x) := [\sigma(x)w]$ for $x \in M(X)$.

(b) *The (m) -cohomology induced map of X :*

$$\psi_m(X) : H_m(\text{Der} M(X)) \rightarrow \text{Der}_m H^*(X; \mathbb{Q})$$

is given by $\psi_m(X)([\sigma])(w) := [\sigma(w)]$.

Proposition 1.7. *For the odd-sphere S^n ($n > 1$),*

(a) *$\text{Baut}_1 X$ has a weak retraction for $\text{Baut}_1(X \times S^n)$ if $\mu_{m,k}(X) = 0$ for any $0 \leq k < n < m$.*

(b) *$\text{Baut}_1 S^n$ has a weak retraction for $\text{Baut}_1(X \times S^n)$ if $\psi_m(X) = 0$ for any $0 < m < n$.*

Definition 1.8. ([25]) *A simply connected space X with Sullivan minimal model $M(X) = (\Lambda V, d)$ is said to be formal if there is a quasi-isomorphism $M(X) \rightarrow (H^*(X; \mathbb{Q}), 0)$ and X is said to be coformal if there is a quasi-isomorphism from the Quillen's DGL-model [17] of X to the rational homotopy Lie algebra $\pi_*(\Omega X)_{\mathbb{Q}}$ (equivalently, the differential of Sullivan minimal model is quadratic, i.e., $dV \subset \Lambda^2 V$).*

When $\text{Baut}_1(X \times S^n)$ is coformal, “ $\text{Baut}_1 X$ has a rational retraction for $\text{Baut}_1(X \times S^n)$ ” is equivalent to “ $\text{Baut}_1 X$ has a weak retraction for $\text{Baut}_1(X \times S^n)$ ” from Lemma 2.2 below.

Theorem 1.9. *Suppose that $H^*(\text{Baut}_1 X; \mathbb{Q})$ is free. Then the differential of $M(\text{Baut}_1(X \times S^n))$ does not have a quadratic part for an odd-sphere S^n if (a) $\mu_{m,k}(X) = 0$ for any $0 \leq k < n < m$ and (b) $\psi_m(X) = 0$ for any $0 < m < n$.*

In particular, when $\text{Baut}_1(X \times S^n)$ is coformal, $H^*(\text{Baut}_1(X \times S^n); \mathbb{Q})$ is free if (a) and (b) hold.

An elliptic space X is said to be *pure* if $dV^{\text{even}} = 0$ and $dV^{\text{odd}} \subset \Lambda V^{\text{even}}$ for $M(X) = (\Lambda V, d)$. For example, simply connected Lie groups, homogeneous spaces and F_0 -spaces are pure spaces. More generally, X is said to be *two-stage* if $dV_0 = 0$ and $dV_1 \subset \Lambda V_0$ for $V = V_0 \oplus V_1$. From degree arguments, we get

Example 1.10. (1) *$H^*(\text{Baut}_1 X; \mathbb{Q})$ is free if and only if $H^*(\text{Baut}_1(X \times S^2); \mathbb{Q})$ is free.*

(2) *When X is elliptic, $H^*(\text{Baut}_1(X \times S^3); \mathbb{Q})$ is free if and only if $M(X) \cong (\Lambda(x_1, \dots, x_m, y_1, \dots, y_n), d)$ with $|x_1| = \dots = |x_m| = 2$ and $|y_1| = \dots = |y_n| = 3$.*

In the followings, the symbol (v, f) means the *elementary derivation* that takes a generator v of V to an element f of ΛV and the other generators to 0.

Example 1.11. Let X be the pullback of the sphere bundle of the tangent bundle of S^{a+b} by the canonical degree 1 map $S^a \times S^b \rightarrow S^{a+b}$. Then it is the total space of a fibration $S^{a+b-1} \rightarrow X \rightarrow S^a \times S^b$. Notice that $M(X) = (\Lambda(v_1, v_2, v_3), d)$ where $|v_1| = a$, $|v_2| = b$, $|v_3| = a + b - 1$ are odd, $dv_1 = dv_2 = 0$ and $dv_3 = v_1 v_2$. Then

$$H^*(X; \mathbb{Q}) \cong \frac{\Lambda(v_1, v_2) \otimes \mathbb{Q}[w_1, w_2]}{(v_1 v_2, v_1 w_1, v_2 w_2, v_1 w_2 + v_2 w_1, w_1^2, w_2^2, w_1 w_2)}$$

with $w_1 = [v_1 v_3]$ and $w_2 = [v_2 v_3]$, so X is not formal. From §3(II), $H^*(Baut_1 X; \mathbb{Q})$ is an one or two variable free algebra: $\Lambda(v_{3,0})$ or $\Lambda(v_{2,1}, v_{3,0})$. By referring to Theorem 1.9, a non-freeness condition of $H^*(Baut_1(X \times S^n); \mathbb{Q})$ for an odd integer n is given by (a) $\mu_{|v_3|,0}([(v_3, 1)], 1) = [(v_3, 1)]$, $\mu_{|v_2|-|v_1|,0}([(v_2, v_1)], 1) = [(v_2, v_1)]$ or (b) $\psi_{|v_3|}([(v_3, 1)]) = (w_1, v_1) + (w_2, v_2)$ under certain degree conditions. From Theorem 4.1 below, we have that $H^*(Baut_1(X \times S^n); \mathbb{Q})$ is free if and only if $a = b$ and $2a - 1 \leq n < 3a - 1$. In §4, we give the Sullivan minimal models for all cases of degrees.

In this paper, second, we consider about Question 1.1 for certain pure spaces X .

Theorem 1.12. Let X be a non-formal pure space where $M(X) = (\Lambda(x_1, x_2, y_1, y_2, y_3), d)$ where $|x_i|$ is even with $|x_1| \leq |x_2|$ and $|y_i|$ is odd with $|y_1| \leq |y_2| \leq |y_3|$. Let $dy_i = f_i \in \mathbb{Q}[x_1, x_2]$ for $i = 1, 2, 3$. Then $H^*(Baut_1 X; \mathbb{Q})$ is not a polynomial algebra if and only if the following (I) or (II) holds:

(I) There is an odd-element $w \in H^*(X; \mathbb{Q})$ with $|w| < |y_3|$.

(II) When $|x_1| < |x_2|$,

$$f_1 = x_1^l \in \mathbb{Q}[x_1], \quad \partial f_2 / \partial x_2 \cdot x_1^k \in (f_1), \quad \partial f_3 / \partial x_2 \cdot x_1^k \in (f_1, f_2) \quad (*)$$

for some $1 < l$ and $0 < k < \min\{l, |x_2|/|x_1|\}$. Here $\partial f_i / \partial x_j$ is the partial differentiation of f_i by x_j and (S) is the ideal of $\mathbb{Q}[x_1, x_2]$ generated by a set S .

In the following examples, (1) and (2) correspond to (I) and (II) of the above theorem, respectively. Furthermore (3) satisfies the both cases of (I) and (II). In all cases, $Baut_1 X$ are not formal.

Example 1.13. (1) When X is the total space of a fibration $S^3 \rightarrow X \rightarrow S^2 \times \mathbb{CP}^3$ such that $M(X) = (\Lambda(x_1, x_2, y_1, y_2, y_3), d)$ with $|x_1| = |x_2| = 2$, $|y_1| = |y_2| = 3$, $|y_3| = 7$, $dy_1 = x_1^2$, $dy_2 = x_1 x_2$, $dy_3 = x_2^4$. There is the even degree non-zero element $\tau = [(y_3, x_2 y_1 - x_1 y_2)] \in H_2(DerM(X))$. Then we have

$$M(Baut_1 X) \cong (\Lambda(v_2, v'_2, v''_2, v_3, v_4, v'_4, v''_4, v_6, v'_6, v_8), d)$$

with $|v_i| = |v'_i| = |v''_i| = i$, $dv_2 = dv'_2 = dv''_2 = dv_3 = dv_4 = dv'_4 = dv_8 = 0$, $dv''_4 = v_2 v_3$, $dv_6 = v_3 v_4$ and $dv'_6 = v_3 v'_4$. Here the element v_3 corresponds to σ (with respect to Theorem 2.1 below).

(2) When X is the homogeneous space $SU(6)/SU(3) \times SU(3)$, $M(X) = (\Lambda(x_1, x_2, y_1, y_2, y_3), d)$ with $|x_1| = 4$, $|x_2| = 6$, $|y_1| = 7$, $|y_2| = 9$, $|y_3| = 11$, $dx_1 = dx_2 = 0$, $dy_1 = x_1^2$, $dy_2 = x_1 x_2$, $dy_3 = x_2^2$ [8]([1]). There is the even degree non-zero element $\sigma = [(x_2, x_1) + (y_2, y_1) + 2(y_3, y_2)] \in H_2(DerM(X))$. Then we have

$$M(Baut_1 X) \cong (\Lambda(v_2, v_3, v_4, v_6, v_8, v'_8, v_{10}, v_{12}), d)$$

with $|v_i| = |v'_i| = i$, $dv_2 = dv_3 = dv_8 = 0$, $dv_4 = v_2v_3$, $dv_6 = v_3v_4$, $dv'_8 = v_3v_6$, $dv_{10} = v_3v_8$ and $dv_{12} = v_3v_{10}$. Here the element v_3 corresponds to σ .

(3) When X is the total space of a fibration $S^5 \rightarrow X \rightarrow S^2 \times \mathbb{H}P^2$ such that $M(X) = (\Lambda(x_1, x_2, y_1, y_2, y_3), d)$ with $|x_1| = 2$, $|x_2| = 4$, $|y_1| = 3$, $|y_2| = 5$, $|y_3| = 11$, $dy_1 = x_1^2$, $dy_2 = x_1x_2$, $dy_3 = x_2^2$. There are the even degree non-zero elements $\sigma = [(x_2, x_1) + (y_2, y_1) + 3(y_3, x_2y_2)] \in H_2(\text{Der}M(X))$ and $\tau = [(y_3, x_2y_1 - x_1y_2)] \in H_4(\text{Der}M(X))$. Then we have

$$M(\text{Baut}_1X) \cong (\Lambda(v_2, v_3, v_4, v'_4, v_5, v_6, v_8, v_{10}, v_{12}), d)$$

with $|v_i| = |v'_i| = i$, $dv_2 = dv_3 = dv_4 = dv_5 = 0$, $dv'_4 = v_2v_3$, $dv_6 = v_3v_4$, $dv_8 = v_2v_3v'_4 + v_4v_5 + v_3v_6$, $dv_{10} = v_3v_8 + v_5v_6$ and $dv_{12} = v_3v'_4v_6$. Here the elements v_3 and v_5 correspond to σ and τ , respectively. It is not even coformal.

Finally we give an example of a non-formal homogeneous space X that does not satisfy the conditions of Theorem 1.12.

Example 1.14. Let X be a homogeneous space with $X = SU(2) \times SU(2) \times SU(2)/T^2$ of $T^2 = S^1 \times S^1$. Due to [16], there are two types of torus embeddings $T^2 \subset SU(2) \times SU(2) \times SU(2)$ such that $M(X) = (\Lambda(x_1, x_2, y_1, y_2, y_3), d)$ with $|x_1| = |x_2| = 2$, $|y_1| = |y_2| = |y_3| = 3$ and

- (i) $dx_1 = dx_2 = 0$, $dy_1 = x_1^2$, $dy_2 = x_2^2$, $dy_3 = 0$ (formal)
- (ii) $dx_1 = dx_2 = 0$, $dy_1 = x_1^2$, $dy_2 = x_2^2$, $dy_3 = x_1x_2$ (non-formal).

In both cases,

$$H^*(\text{Baut}_1X; \mathbb{Q}) \cong \mathbb{Q}[u_1, u_2, u_3, u_4, u_5, u_6, u_7]$$

with $|u_1| = |u_2| = |u_3| = |u_4| = 2$ and $|u_5| = |u_6| = |u_7| = 4$ since $\pi_{*+1}(\text{Baut}_1X)_{\mathbb{Q}} \cong H_*(\text{Der}M(X)) \cong \mathbb{Q}\{(y_1, x_2), (y_2, x_k), (y_3, x_1), (y_3, x_2), (y_1, 1), (y_2, 1), (y_3, 1)\}$ in which $k = 1$ for (i) and $k = 2$ for (ii) from Theorem 2.1.

In §2, we prepare about computation of derivations and give the proofs of Theorems 1.9 and 1.12. In §3, we compare $M(\text{Baut}_1X)$ for formal cases and non-formal cases on elliptic spaces X with rational homotopy groups of rank 3. In §4, we give the Sullivan minimal models on all cases of Example 1.11. In §5, we give the explicit computation for giving the minimal model of Example 1.13 (2) but omit the computations of Example 1.13 (1) and (3) since they are similar to it.

2. SULLIVAN MODELS AND DERIVATIONS

We use the *Sullivan minimal model* $M(X)$ of a simply connected space X of finite type. It is a free $\mathbb{Q}$ -commutative differential graded algebra (abbr., DGA) $(\Lambda V, d)$ with a $\mathbb{Q}$ -graded vector space $V = \bigoplus_{i>1} V^i$ where $\dim V^i < \infty$ and a decomposable differential, i.e.,

$$d(V^i) \subset (\Lambda^+V \cdot \Lambda^+V)^{i+1} \text{ and } d \circ d = 0.$$

Here Λ^+V is the ideal of ΛV generated by elements of positive degree. We often denote $(\Lambda V, d)$ simply by ΛV . The degree of an element x of a graded algebra is denoted by $|x|$. Then we have

$$xy = (-1)^{|x||y|}yx \text{ and } d(xy) = d(x)y + (-1)^{|x|}x d(y).$$

We note that $M(X)$ determines the rational homotopy type $X_{\mathbb{Q}}$ of X . In particular, there are the following isomorphisms

$$\text{Hom}(V^i, \mathbb{Q}) \cong \pi_i(X) \otimes \mathbb{Q} (= \pi_i(X)_{\mathbb{Q}}) \text{ and } H^*(\Lambda V, d) \cong H^*(X; \mathbb{Q})$$

See [3] for a general introduction and the standard notations.

Let $\text{Der}_i M(X)$ be the set of $\mathbb{Q}$ -derivations of $M(X)$ decreasing the degree by i with $\sigma(xy) = \sigma(x)y + (-1)^{i|x|}x\sigma(y)$ for $x, y \in M(X)$. The boundary operator $\partial : \text{Der}_i M(X) \rightarrow \text{Der}_{i-1} M(X)$ is defined by $\partial(\sigma) = d \circ \sigma - (-1)^i \sigma \circ d$ for $\sigma \in \text{Der}_i M(X)$. We denote $\bigoplus_{i>0} \text{Der}_i M(X)$ by $\text{Der} M(X)$ in which $\text{Der}_1 M(X)$ is ∂ -cycles. Then $\text{Der} M(X)$ is a DGL by the Lie bracket $[\sigma, \tau] := \sigma \circ \tau - (-1)^{|\sigma||\tau|} \tau \circ \sigma$. Furthermore, recall the definition (sign convention) of D.Tannré [25, p.25]: $C^*(L, \partial) = (\Lambda s^{-1} \sharp L, D = d_1 + d_2)$ with

- (i) $\langle d_1 s^{-1} z; sx \rangle = -\langle z; \partial x \rangle$ and
- (ii) $\langle d_2 s^{-1} z; sx_1, sx_2 \rangle = (-1)^{|x_1|} \langle z; [x_1, x_2] \rangle,$

where $\langle s^{-1} z; sx \rangle = (-1)^{|z|} \langle z; x \rangle$ for a DGL (L, ∂) and $\sharp L$ is the dual space of L .

Theorem 2.1. ([24, p.314], [4]) *The DGA $C^*(\text{Der} M(X), \partial)$ is the Sullivan model $m(\text{Baut}_1 X)$ of $\text{Baut}_1 X$. In particular, $\pi_*(\Omega \text{Baut}_1 X) \otimes \mathbb{Q} \cong H_*(\text{Der} M(X), \partial)$ as graded Lie algebras, in which the left hand has the Samelson bracket.*

Here the DGA $m(\text{Baut}_1 X)$ need not be minimal (cf. Proposition 5.2 below). The following lemma is obvious from DGA-arguments.

Lemma 2.2. *Suppose that a map $f : S \rightarrow T$ is $\pi_*(f)_{\mathbb{Q}}$ -injective. Then the model is given as $M(f) : M(T) = (\Lambda(U \oplus V), D) \rightarrow (\Lambda V, d) = M(S)$ with*

$$M(f)|_V = \text{id}_V, \quad M(f)(U) = 0, \quad d = \overline{D}, \quad DU \in \Lambda^+ U \otimes \Lambda V, \quad DV \in \Lambda U \otimes \Lambda^+ V.$$

Then S is a rational factor of T if and only if $DU \subset \Lambda U$ and $D|_V = d$. Furthermore

- (1) *S has a rational retraction for T if and only if $DV \subset \Lambda V$ (i.e., $D|_V = d$).*
- (2) *S has a weak retraction for T if and only if $DV \subset \Lambda V \oplus \Lambda^{>2}(U \oplus V)$.*

Proposition 2.3. *Let $M(X \times S^n) = (\Lambda(V \oplus u), d)$ where $M(X) = (\Lambda V, d)$ and $|u| = n$ for an odd-integer n . Let $A = \bigoplus_{i>0} A_i$, $B = \bigoplus_{i>0} B_i$ and $C = \bigoplus_{i>0} C_i$ where*

$$H_i(\text{Der}(\Lambda(V \oplus u))) = A_i \oplus B_i \oplus C_i$$

$$:= H_i(\text{Der}(\Lambda V)) \oplus H_i(\text{Der}(\Lambda V, (u))) \oplus H_i(\text{Der}(\Lambda(u), \Lambda V)),$$

where (u) is the ideal of $\Lambda(V \oplus u)$ generated by u . Then

- (1) *$\text{Baut}_1 X$ has a weak retraction for $\text{Baut}_1(X \times S^n)$ if $[B, C] = 0$.*
- (2) *$\text{Baut}_1 S^n$ has a weak retraction for $\text{Baut}_1(X \times S^n)$ if $[A, C] = 0$.*
- (3) *$\pi_*(\Omega \text{Baut}_1(X \times S^n))_{\mathbb{Q}}$ is abelian (i.e., the Lie bracket of $H_*(\text{Der}(\Lambda(V \oplus u)))$ is trivial) if $[A, A] = 0$, $[A, C] = 0$ and $[B, C] = 0$.*

Proof. In general, since they are negative degree derivations, we have

$$[A, A] = A, \quad [A, B] = B, \quad [A, C] = C, \quad [B, B] = B, \quad [B, C] = A, \quad [C, C] = 0.$$

From Theorem 2.1 and Lemma 2.2, we have (1),(2) and (3). In particular, if $[A, A] = 0$, we have $[A, B] = [B, B] = 0$. Thus (3) holds. $\square$

Claim 2.4. (1) $\mu_{m,k}(X) = 0$ for any $0 \leq k < n < m$ if and only if $[B, C] = 0$.

(2) $\psi_m(X) = 0$ for any $0 < m < n$ if and only if $[A, C] = 0$.

Proof. (1) For $\sigma \in H_m(\text{Der}M(X))$ and $w \in H^k(X; \mathbb{Q})$,

$$\mu_{m,k}(\sigma \otimes w) = \sigma \otimes w = \pm[\sigma \otimes u, (u, w)] \in [B, C]$$

under $0 \leq k < n < m$. Conversely, all elements of $[B, C]$ are given as them.

(2) For $\sigma \in H_m(\text{Der}M(X))$ and $w \in H^k(X; \mathbb{Q})$,

$$(u, \psi_m(\sigma)(w)) = (u, \sigma(w)) = \pm[\sigma, (u, w)] \in [A, C]$$

under $0 < m \leq k < n$. Conversely, all elements of $[A, C]$ are given as them. $\square$

Proof of Proposition 1.7. (a) It holds from Proposition 2.3 (1) and Claim 2.4 (1).

(b) It holds from Proposition 2.3 (2) and Claim 2.4 (2). $\square$

Proof of Theorem 1.9. It holds from Proposition 2.3 (3) and Claim 2.4 (1)(2). $\square$

From Theorem 2.1, $\pi_{*+1}(\text{Baut}_1 Y) \otimes \mathbb{Q} \cong H_*(\text{Der}M(Y))$. Thus we have immediately by degree arguments

Lemma 2.5. $H_{\text{even}}(\text{Der}M(X)) = 0$ if and only if $H^*(\text{Baut}_1 X; \mathbb{Q})$ is a polynomial algebra.

Proof of Theorem 1.12. From Lemma 2.5, it is sufficient to observe when a non-zero even degree element σ exists in $H_*(\text{Der}M(X))$. Then we can divide it two cases as (I) : “ $|v|$ and $|f|$ are both odd” or (II) : “ $|v|$ and $|f|$ are both even” for $\sigma = (v, f) + \dots$.

A possibility of $H_{\text{even}}(\text{Der}M(X)) \neq 0$ is that there is the non-exact δ -cycle $\sigma = (y_3, w)$ for some odd-degree non-exact cocycle w in the ideal (y_1, y_2) , which is equivalent to (I).

The other possibility is given, when $|x_1| < |x_2|$, as that there exists a non-exact δ -cycle $\sigma = (x_2, x_1^k) + \tau$ for some $0 < k < |x_2|/|x_1|$ and a derivation τ . Let $dy_i = f_i \in \mathbb{Q}[x_1, x_2]$ for $i = 1, 2, 3$. There exists such a τ if and only if there are elements $g, h_1, h_2 \in \mathbb{Q}[x_1, x_2]$ such that

$$\begin{aligned} \partial(x_2, x_1^k) &= -(y_1, \partial f_1 / \partial x_2 \cdot x_1^k) - (y_2, \partial f_2 / \partial x_2 \cdot x_1^k) - (y_3, \partial f_3 / \partial x_2 \cdot x_1^k) \\ &= (y_2, g f_1) + (y_3, h_1 f_1 + h_2 f_2) = \partial((y_2, g y_1) + (y_3, h_1 y_1 + h_2 y_2)) = \partial(-\tau). \end{aligned}$$

It is equivalent to (*) of (II). $\square$

Remark 2.6. In Theorem 1.12(I), if $\psi_{|y_i|}(y_i, 1)(w) = [\partial w / \partial y_i] \neq 0$, there is the non-exact δ -cycle $\sigma = (y_3, \partial w / \partial y_i)$ with $\sigma = [(y_i, 1), (y_3, w)]$ for non-exact δ -cycles $(y_i, 1)$ and (y_3, w) ($i = 1$ or 2). Then the rational homotopy Lie algebra $\pi_*(\Omega \text{Baut}_1 X)_{\mathbb{Q}}$ has a non-trivial Lie bracket (is not abelian) from Theorem 2.1. Especially $H^*(\text{Baut}_1 X; \mathbb{Q})$ is not even free.

3. ODDLY GENERATED MODELS OF RANK 3

For elliptic spaces X with $\dim \pi_*(X)_{\mathbb{Q}} = 3$ and $\pi_*(X)_{\mathbb{Q}} = \pi_{\text{odd}}(X)_{\mathbb{Q}}$, let $M(X) = (\Lambda(v_1, v_2, v_3), d)$ with $|v_i|$ odd for $i = 1, 2, 3$ of $|v_1| \leq |v_2| \leq |v_3|$. Then the minimal model $M := M(\text{Baut}_1 X)$ is given as follows:

(I) *Formal case:* $d(v_1) = d(v_2) = d(v_3) = 0$.

$$(1.1) \quad |v_1| = |v_2| = |v_3|$$

$$M \cong (\Lambda(v_{1,0}, v_{2,0}, v_{3,0}), 0)$$

$$(1.2) \quad |v_1| = |v_2| < |v_3| \leq 2|v_1|$$

$$M \cong (\Lambda(v_{1,0}, v_{2,0}, v_{3,0}, v_{3,1}, v_{3,2}), D)$$

with $D(v_{1,0}) = D(v_{2,0}) = D(v_{3,1}) = D(v_{3,2}) = 0$ and $D(v_{3,0}) = v_{1,0}v_{3,1} + v_{2,0}v_{3,2}$.

$$(1.3) \quad |v_1| = |v_2| < \frac{1}{2}|v_3|$$

$$M \cong (\Lambda(v_{1,0}, v_{2,0}, v_{3,0}, v_{3,1}, v_{3,2}, v_{3,12}), D)$$

with $D(v_{1,0}) = D(v_{2,0}) = D(v_{3,12}) = 0$, $D(v_{3,1}) = v_{1,0}v_{3,12}$, $D(v_{3,2}) = v_{2,0}v_{3,12}$ and $D(v_{3,0}) = v_{1,0}v_{3,1} + v_{2,0}v_{3,2}$.

$$(1.4) \quad |v_1| < |v_2| < |v_3| \leq |v_1| + |v_2|$$

$$M \cong (\Lambda(v_{1,0}, v_{2,0}, v_{2,1}, v_{3,0}, v_{3,1}, v_{3,2}), D)$$

with $D(v_{1,0}) = D(v_{2,1}) = 0$, $D(v_{2,0}) = v_{1,0}v_{2,1}$, $D(v_{3,1}) = v_{2,1}v_{3,2}$ and $D(v_{3,0}) = v_{1,0}v_{3,1} + v_{2,0}v_{3,2}$.

$$(1.5) \quad |v_1| < |v_2| \text{ and } |v_1| + |v_2| < |v_3|$$

$$M \cong (\Lambda(v_{1,0}, v_{2,0}, v_{2,1}, v_{3,0}, v_{3,1}, v_{3,2}, v_{3,12}), D)$$

with $D(v_{1,0}) = D(v_{2,1}) = D(v_{3,2}) = D(v_{3,12}) = 0$, $D(v_{2,0}) = v_{1,0}v_{2,1}$, $D(v_{3,1}) = v_{2,1}v_{3,2} + v_{3,12}v_{2,0}$ and $D(v_{3,0}) = v_{1,0}v_{3,1} + v_{2,0}v_{3,2}$.

$$(1.6) \quad |v_1| < |v_2| = |v_3|$$

$$M \cong (\Lambda(v_{1,0}, v_{2,0}, v_{2,1}, v_{3,0}, v_{3,1}), D)$$

with $D(v_{1,0}) = D(v_{2,1}) = D(v_{3,1}) = 0$, $D(v_{2,0}) = v_{1,0}v_{2,1}$ and $D(v_{3,0}) = v_{1,0}v_{3,1}$.

(II) *Non-formal case:* $d(v_1) = d(v_2) = 0$ and $d(v_3) = v_1v_2$.

$$(2.1) \quad |v_1| = |v_2|$$

$$M \cong (\Lambda(v_{3,0}), 0).$$

$$(2.2) \quad |v_1| < |v_2|$$

$$M \cong (\Lambda(v_{2,1}, v_{3,0}), 0).$$

Here the elements $v_{i,j}$, $v_{i,jk}$ and $v_{i,0}$ corresponds to the derivations (v_i, v_j) , (v_i, v_jv_k) and $(v_i, 1) = v_i^*$, respectively with respect to Theorem 2.1. Thus $|v_{i,j}| = |v_i| - |v_j| + 1$ is odd and $|v_{i,jk}| = |v_i| - |v_j| - |v_k| + 1$, $|v_{i,0}| = |v_i| + 1$ are even. We also adopt these symbols to the following sections again.

4. EXAMPLE 1.11

Let's directly compute the rational homotopies of $Baut_1(X \times S^n)$ when $M(X) = (\Lambda(v_1, v_2, v_3), d)$ with $d(v_1) = d(v_2) = 0$ and $d(v_3) = v_1v_2$ (i.e., (II) of §3) and $H^*(S^n; \mathbb{Q}) = \Lambda(u)$ with $|u| = n$ odd. The following cases (1) $\sim$ (6) exist:

$$(1) \quad |u| \leq |v_1|$$

$$(i) \quad |u| + |v_1| < |v_2|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{u,0}, v_{2,1}, v_{2,u1}, v_{3,0}, v_{3,u}), d)$$

with $d(v_{3,0}) = v_{u,0}v_{3,u}$ and $d(v_{2,1}) = v_{u,0}v_{2,u1}$ ($dv = 0$ for the other generators v).

$$(ii) \quad |u| + |v_1| > |v_2| \text{ and } |v_1| < |v_2|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{u,0}, v_{2,1}, v_{3,0}, v_{3,u}), d)$$

with $d(v_{3,0}) = v_{u,0}v_{3,u}$.

$$(iii) \quad |v_1| = |v_2|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{u,0}, v_{3,0}, v_{3,u}), d)$$

with $d(v_{3,0}) = v_{u,0}v_{3,u}$.

$$(2) \quad |v_1| < |u| \leq |v_2|$$

$$(i) \quad |u| + |v_1| < |v_2|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{u,0}, v_{u,1}, v_{2,1}, v_{2,u1}, v_{3,0}, v_{3,u}), d)$$

with $d(v_{3,0}) = v_{u,0}v_{3,u}$ and $d(v_{2,1}) = v_{u,0}v_{2,u1}$.

$$(ii) \quad |u| + |v_1| > |v_2|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{u,0}, v_{u,1}, v_{2,1}, v_{3,0}, v_{3,u}), d)$$

with $d(v_{3,0}) = v_{u,0}v_{3,u}$.

$$(3) \quad |v_1| < |v_2| < |u| < |v_3|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{2,1}, v_{u,0}, v_{u,1}, v_{u,2}, v_{3,0}, v_{3,u}), d)$$

with $d(v_{3,0}) = v_{u,0}v_{3,u}$, $d(v_{u,0}) = v_{u,1}v_{u,2}v_{3,u}$ and $d(v_{u,1}) = v_{2,1}v_{u,2}$.

$$(4) \quad |v_1| = |v_2| < |u| < |v_3|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{u,0}, v_{u,1}, v_{u,2}, v_{3,0}, v_{3,u}), d)$$

with $d(v_{3,0}) = v_{u,0}v_{3,u}$ and $d(v_{u,0}) = v_{u,1}v_{u,2}v_{3,u}$.

$$(5) \quad |v_1| < |v_2| < |v_3| \leq |u|$$

$$(i) \quad |v_1| + |v_2| + |v_3| < |u|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{2,1}, v_{3,0}, v_{u,0}, v_{u,1}, v_{u,2}, v_{u,13}, v_{u,23}, v_{u,123}), d)$$

with $d(v_{u,0}) = v_{3,0}^2 v_{u,123}$, $d(v_{u,1}) = v_{2,1}v_{u,2} + v_{3,0}v_{u,13}$, $d(v_{u,2}) = v_{3,0}v_{u,23}$ and $d(v_{u,13}) = v_{2,1}v_{u,23}$.

$$(ii) \quad |v_2| + |v_3| < |u| \leq |v_1| + |v_2| + |v_3|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{2,1}, v_{3,0}, v_{u,0}, v_{u,1}, v_{u,2}, v_{u,13}, v_{u,23}), d)$$

with $d(v_{u,1}) = v_{2,1}v_{u,2} + v_{3,0}v_{u,13}$, $d(v_{u,2}) = v_{3,0}v_{u,23}$ and $d(v_{u,13}) = v_{2,1}v_{u,23}$.

$$(iii) \quad |v_1| + |v_3| < |u| < |v_2| + |v_3|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{2,1}, v_{3,0}, v_{u,0}, v_{u,1}, v_{u,2}, v_{u,13}), d)$$

with $d(v_{u,1}) = v_{2,1}v_{u,2} + v_{3,0}v_{u,13}$.

$$(iv) |u| < |v_1| + |v_3|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{2,1}, v_{3,0}, v_{u,0}, v_{u,1}, v_{u,2}), d)$$

with $d(v_{u,1}) = v_{2,1}v_{u,2}$.

$$(6) |v_1| = |v_2| < |v_3| \leq |u|$$

$$(i) |v_1| + |v_2| + |v_3| < |u|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{3,0}, v_{u,0}, v_{u,1}, v_{u,2}, v_{u,13}, v_{u,23}, v_{u,123}), d)$$

with $d(v_{u,1}) = v_{3,0}v_{u,13}$ and $d(v_{u,2}) = v_{3,0}v_{u,23}$.

$$(ii) |v_2| + |v_3| < |u| \leq |v_1| + |v_2| + |v_3|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{3,0}, v_{u,0}, v_{u,1}, v_{u,2}, v_{u,13}, v_{u,23}), d)$$

with $d(v_{u,1}) = v_{3,0}v_{u,13}$ and $d(v_{u,2}) = v_{3,0}v_{u,23}$.

$$(iii) |u| < |v_1| + |v_3|$$

$$M(Baut_1(X \times S^n)) \cong (\Lambda(v_{3,0}, v_{u,0}, v_{u,1}, v_{u,2}), 0).$$

Here $v_{u,0}$, $v_{u,i}$, $v_{i,u}$, $\dots$ correspond to the derivations $(u, 1)$, (u, v_i) , (v_i, u) , $\dots$ respectively. Thus $|v_{u,0}|$, $|v_{u,ij}|$, $|v_{i,u}|$ are even and $|v_{i,u}|$, $|v_{u,i}|$, $|v_{u,ijk}|$ are odd.

Thus we have the following

Theorem 4.1. *When $M(X) = (\Lambda(v_1, v_2, v_3), d)$ with $|v_i|$ odd, $dv_1 = dv_2 = 0$ and $dv_3 = v_1v_2$, there are 14-types of $Baut_1(X \times S^n)$ with n odd as*

type	f_X	r_X	$w.r_X$	f_{S^n}	r_{S^n}	$w.r_{S^n}$	formal	coformal	rank	H^* -free
$(1)_i$	no	no	no	no	yes	yes	no	yes	5	no
$(1)_{ii}$	no	no	no	no	yes	yes	no	yes	4	no
$(1)_{iii}$	no	no	no	no	yes	yes	no	yes	3	no
$(2)_i$	no	no	no	no	yes	yes	no	yes	6	no
$(2)_{ii}$	no	no	no	no	yes	yes	no	yes	5	no
(3)	no	no	no	no	no	yes	no	no	6	no
(4)	no	no	no	no	no	yes	no	no	5	no
$(5)_i$	no	yes	yes	no	no	yes	no	no	8	no
$(5)_{ii}$	no	yes	yes	yes	yes	yes	no	yes	7	no
$(5)_{iii}$	no	yes	yes	yes	yes	yes	yes	yes	6	no
$(5)_{iv}$	no	yes	yes	yes	yes	yes	no	yes	5	no
$(6)_i$	no	yes	yes	yes	yes	yes	yes	yes	7	no
$(6)_{ii}$	no	yes	yes	yes	yes	yes	yes	yes	6	no
$(6)_{iii}$	yes	yes	yes	yes	yes	yes	yes	yes	4	yes

Here f_X means that $Baut_1X$ is a rational factor of $Baut_1(X \times S^n)$, r_X means that $Baut_1X$ has a rational retraction for $Baut_1(X \times S^n)$ and $w.r_X$ means that $Baut_1X$ has a weak retraction for $Baut_1(X \times S^n)$. Furthermore f_{S^n} , r_{S^n} and $w.r_{S^n}$ are similar symbols for $Baut_1S^n$. H^* -free means that $H^*(Baut_1(X \times S^n); \mathbb{Q})$ is free.

Proof. Since $M(\text{Baut}_1 X) = (\Lambda v_{3,0}, 0)$ or $(\Lambda(v_{3,0}, v_{2,1}), 0)$ and $M(\text{Baut}_1 S^n) = (\Lambda v_{u,0}, 0)$, we can check whether or not the properties $f_X, \dots, w.r_{S^n}$ are true respectively from Lemma 2.2. See [21, §4] for formality conditions. $\square$

Corollary 4.2. *Suppose $M(X) = (\Lambda(v_1, v_2, v_3), d)$ where $|v_i|$ are odd, $dv_1 = dv_2 = 0$, $dv_3 = v_1 v_2$ and n is odd. Then*

(1) *The followings are equivalent:*

(i) *$H^*(\text{Baut}_1(X \times S^n); \mathbb{Q})$ is free,*

(ii) *$|v_1| = |v_2| < |v_3| \leq n < |v_1| + |v_3|$ and*

(iii) *$\text{Baut}_1 X$ is a rational factor of $\text{Baut}_1(X \times S^n)$.*

(2) *$\text{Baut}_1(X \times S^n)$ is coformal if and only if $\text{Baut}_1 S^n$ has a rational retraction for $\text{Baut}_1(X \times S^n)$.*

(3) *If $\text{Baut}_1(X \times S^n)$ is formal, $\text{Baut}_1 S^n$ is a rational factor of $\text{Baut}_1(X \times S^n)$ and $\text{Baut}_1 X$ has a rational retraction for $\text{Baut}_1(X \times S^n)$.*

(4) *$\text{Baut}_1 X$ has a rational retraction for $\text{Baut}_1(X \times S^n)$ if and only if it does a weak retraction.*

(5) *$\text{Baut}_1 S^n$ has a weak retraction for $\text{Baut}_1(X \times S^n)$.*

5. CASE OF $X = SU(6)/SU(3) \times SU(3)$

Let $X = SU(6)/SU(3) \times SU(3)$. Then $M(X) = (\Lambda(x_1, x_2, y_1, y_2, y_3), d)$ with $|x_1| = 4$, $|x_2| = 6$, $|y_1| = 7$, $|y_2| = 9$, $|y_3| = 11$, $dx_1 = dx_2 = 0$, $dy_1 = x_1^2$, $dy_2 = x_1 x_2$ and $dy_3 = x_2^2$. Then a basis of $\text{Der}M(X)$ is given as the table:

degree	generators
11	$(y_3, 1)$
9	$(y_2, 1)$
7	$(y_3, x_1), (y_1, 1)$
6	$(x_2, 1)$
5	$(y_3, x_2), (y_2, x_1)$
4	$(y_3, x_1), (x_1, 1)$
3	$(y_3, x_1^2), (y_2, x_2), (y_1, x_1)$
2	$(y_3, x_2), (y_2, x_1), (x_2, x_1)$
1	$(y_3, x_1 x_2), (y_2, x_1^2), (y_1, x_2)$

Proposition 5.1. *For $M(X) = (\Lambda(x_1, x_2, y_1, y_2, y_3), d)$ of above, we have*

$$\pi_{*+1}(\text{Baut}_1 X) \otimes \mathbb{Q} \cong \mathbb{Q}\{s(y_3, 1), s(y_2, 1), s(y_1, 1), s(y_1, x_1), s(y_1, x_2), s(y_3, x_1), s(y_3, x_2), \\ s(2(y_3, y_2) + (y_2, y_1) + (x_2, x_1))\}$$

as a graded vector space. Here $|s((v, f) + \dots)| = |v| - |f| + 1$.

Proof. Recall §2. The differential ∂ on the generators of $\text{Der}M(X)$ is given as

$$\begin{aligned} \partial((y_3, 1)) &= \partial((y_2, 1)) = \partial((y_3, x_1)) = \partial((y_1, 1)) = \partial((y_3, x_2)) = \partial((y_2, x_1)) = 0 \\ \partial((y_3, x_1 x_2)) &= \partial((y_2, x_1^2)) = \partial((y_1, x_2)) = \partial((y_3, x_1^2)) = \partial((y_2, x_2)) = \partial((y_1, x_1)) = 0 \\ \partial((x_2, 1)) &= -(y_2, x_1) - 2(y_3, x_2), \partial((y_3, y_1)) = (y_3, x_1^2), \partial((x_1, 1)) = -2(y_1, x_1) - (y_2, x_2) \\ \partial((y_3, y_2)) &= (y_3, x_1 x_2), \partial((y_2, y_1)) = (y_2, x_1^2), \partial((x_2, x_1)) = -(y_2, x_1^2) - 2(y_3, x_1 x_2). \end{aligned}$$

Then we obtain the result from $\pi_{*+1}(\text{Baut}_1 X) \otimes \mathbb{Q} \cong H_*(\text{Der}M(X)) = \text{Ker}(\partial)/\text{Im}(\partial)$ of Theorem 2.1. $\square$

Proposition 5.2. *The DGA-model $m(\text{Baut}_1 X) := C^*(\text{Der} M(X))$ is given as*

$$m(\text{Baut}_1 X) \cong (\Lambda(V_{y_3,1}, V_{y_2,1}, V_{y_3,x_1}, V_{y_1,1}, V_{x_2,1}, V_{w_1}, V_{y_2,x_1}, V_{y_3,y_1}, V_{x_1,1}, V_{y_3,x_1^2}, V_{y_2,x_2}, V_{w_2}, V_{y_3,y_2}, V_{y_2,y_1}, V_{x_2,x_1}, V_{y_3,x_1x_2}, V_{y_2,x_1^2}, V_{y_1,x_2}), D)$$

with $|V_{y_3,1}| = 12$, $|V_{y_2,1}| = 10$, $|V_{y_3,x_1}| = |V_{y_1,1}| = 8$, $|V_{x_2,1}| = 7$, $|V_{w_1}| = |V_{y_2,x_1}| = 6$, $|V_{y_3,y_1}| = |V_{x_1,1}| = 5$, $|V_{y_3,x_1^2}| = |V_{y_2,x_2}| = |V_{w_2}| = 4$, $|V_{y_3,y_2}| = |V_{y_2,y_1}| = |V_{x_2,x_1}| = 3$, $|V_{y_3,x_1x_2}| = |V_{y_2,x_1^2}| = |V_{y_1,x_2}| = 2$ and differential:

$$\begin{aligned} D(V_{y_3,1}) &= -V_{x_1,1}V_{y_3,x_1} + V_{x_2,1}V_{w_1} - 2V_{x_2,1}V_{y_2,x_1} + V_{y_1,1}V_{y_3,y_1} + V_{y_2,1}V_{y_3,y_2} \\ D(V_{y_2,1}) &= -V_{x_1,1}V_{y_2,x_1} - V_{x_2,1}V_{y_2,x_2} + V_{y_1,1}V_{y_2,y_1} \\ D(V_{y_3,x_1}) &= V_{x_2,x_1}V_{w_1} - 2V_{x_2,x_1}V_{y_2,x_1} - V_{w_2}V_{y_3,y_1} + 2V_{y_2,x_2}V_{y_3,y_1} - 2V_{x_1,1}V_{y_3,x_1^2} \\ &\quad + V_{y_2,x_1}V_{y_3,y_2} - V_{x_2,1}V_{y_3,x_1x_2} \\ D(V_{y_1,1}) &= V_{x_1,1}V_{w_2} - 2V_{x_1,1}V_{y_2,x_2} - V_{x_2,1}V_{y_1,x_2} \\ D(V_{x_2,1}) &= -V_{x_1,1}V_{x_2,x_1} \\ D(V_{w_1}) &= -2V_{x_2,x_1}V_{y_2,x_2} - 4V_{x_1,1}V_{y_2,x_1^2} + 4V_{y_2,x_2}V_{y_2,y_1} - 2V_{w_2}V_{y_2,y_1} - V_{y_1,x_2}V_{y_3,y_1} \\ &\quad - V_{y_2,x_2}V_{y_3,y_2} + V_{x_1,1}V_{y_3,x_1x_2} \\ D(V_{y_2,x_1}) &= V_{x_2,1} - V_{x_2,x_1}V_{y_2,x_2} - V_{w_2}V_{y_2,y_1} + 2V_{y_2,x_2}V_{y_2,y_1} - 2V_{x_1,1}V_{y_2,x_1^2} \\ D(V_{y_3,y_1}) &= -V_{y_2,y_1}V_{y_3,y_2} \\ D(V_{y_3,x_1^2}) &= -V_{y_3,y_1} - V_{x_2,x_1}V_{y_3,x_1x_2} + V_{y_2,x_1^2}V_{y_3,y_2} \\ D(V_{y_2,x_2}) &= V_{x_1,1} + V_{y_1,x_2}V_{y_2,y_1} \\ D(V_{w_2}) &= 2V_{y_1,x_2}V_{y_2,y_1} + V_{x_2,x_1}V_{y_1,x_2} \\ D(V_{y_3,x_1x_2}) &= -V_{y_3,y_2} + 2V_{x_2,x_1} \\ D(V_{y_2,x_1^2}) &= -V_{y_2,y_1} + V_{x_2,x_1} \text{ and} \\ D(V_{x_1,1}) &= D(V_{y_3,y_2}) = D(V_{y_2,y_1}) = D(V_{x_2,x_1}) = D(V_{y_1,x_2}) = 0. \end{aligned}$$

Proof. Recall §2 (i). The differential d_1 is given as

$$\begin{aligned} d_1(V_{y_3,1}) &= d_1(V_{y_2,1}) = d_1(V_{y_3,x_1}) = d_1(V_{y_1,1}) = d_1(V_{x_2,1}) = d_1(V_{y_3,y_1}) = 0, \\ d_1(V_{y_3,y_2}) &= d_1(V_{y_2,y_1}) = d_1(V_{x_2,x_1}) = d_1(V_{x_1,1}) = d_1(V_{y_1,x_2}) = 0, \\ d_1(V_{y_3,x_2}) &= 2V_{x_2,1}, \quad d_1(V_{y_2,x_1}) = V_{x_2,1}, \quad d_1(V_{y_3,x_1^2}) = -V_{y_3,y_1}, \quad d_1(V_{y_2,x_2}) = V_{x_1,1}, \\ d_1(V_{y_1,x_1}) &= 2V_{x_1,1}, \quad d_1(V_{y_3,x_1x_2}) = -V_{y_3,y_2} + 2V_{x_2,x_1}, \quad d_1(V_{y_2,x_1^2}) = -V_{y_2,y_1} + V_{x_2,x_1}. \end{aligned}$$

Next, the Lie bracket is given as

$$\begin{aligned} [(x_1, 1), (y_3, x_1)] &= (y_3, 1), \quad [(x_2, 1), (y_3, x_2)] = (y_3, 1), \quad [(y_1, 1), (y_3, y_1)] = (y_3, 1), \\ [(y_2, 1), (y_3, y_2)] &= (y_3, 1), \quad [(x_1, 1), (y_2, x_1)] = (y_2, 1), \quad [(x_2, 1), (y_2, x_2)] = (y_2, 1), \\ [(y_1, 1), (y_2, y_1)] &= (y_2, 1), \quad [(x_2, x_1), (y_3, x_2)] = (y_3, x_1), \quad [(y_1, x_1), (y_3, y_1)] = (y_3, x_1), \\ \frac{1}{2}[(x_1, 1), (y_3, x_1^2)] &= (y_3, x_1), \quad [(y_2, x_1), (y_3, y_2)] = (y_3, x_1), \quad [(x_2, 1), (y_3, x_1x_2)] = (y_3, x_1), \\ [(x_1, 1), (y_1, x_1)] &= (y_1, 1), \quad [(x_2, 1), (y_1, x_2)] = (y_1, 1), \quad [(x_1, 1), (x_2, x_1)] = (x_2, 1), \\ [(y_1, x_2), (y_3, y_1)] &= (y_3, x_2), \quad [(y_2, x_2), (y_3, y_2)] = (y_3, x_2), \quad [(x_1, 1), (y_3, x_1x_2)] = (y_3, x_2), \\ [(x_2, x_1), (y_2, x_2)] &= (y_2, x_1), \quad [(y_1, x_1), (y_2, y_1)] = (y_2, x_1), \quad \frac{1}{2}[(x_1, 1), (y_2, x_1^2)] = (y_2, x_1), \\ [(y_2, y_1), (y_3, y_2)] &= (y_3, y_1), \quad [(x_2, x_1), (y_3, x_1x_2)] = (y_3, x_1^2), \quad [(y_2, x_1^2), (y_3, y_2)] = (y_3, x_1^2), \end{aligned}$$

$$[(y_1, x_2), (y_2, y_1)] = (y_2, x_2) \quad \text{and} \quad [(x_2, x_1), (y_1, x_2)] = (y_1, x_1).$$

Then the differential d_2 of §2 (ii) is given as

$$\begin{aligned} d_2(V_{y_3,1}) &= -V_{x_1,1}V_{y_3,x_1} - V_{x_2,1}V_{y_3,x_2} + V_{y_1,1}V_{y_3,y_1} + V_{y_2,1}V_{y_3,y_2} \\ d_2(V_{y_2,1}) &= -V_{x_1,1}V_{y_2,x_1} - V_{x_2,1}V_{y_2,x_2} + V_{y_1,1}V_{y_2,y_1} \\ d_2(V_{y_3,x_1}) &= -V_{x_2,x_1}V_{y_3,x_2} + V_{y_1,x_1}V_{y_3,y_1} - 2V_{x_1,1}V_{y_3,x_1^2} + V_{y_2,x_1}V_{y_3,y_2} - V_{x_2,1}V_{y_3,x_1x_2} \\ d_2(V_{y_1,1}) &= -V_{x_1,1}V_{y_1,x_1} - V_{x_2,1}V_{y_1,x_2} \\ d_2(V_{x_2,1}) &= -V_{x_2,1}V_{y_1,x_2} \\ d_2(V_{y_3,x_2}) &= V_{y_1,x_2}V_{y_3,y_1} + V_{y_2,x_2}V_{y_3,y_2} - V_{x_1,1}V_{y_3,x_1x_2} \\ d_2(V_{y_2,x_1}) &= -V_{x_2,x_1}V_{y_2,x_2} + V_{y_1,x_1}V_{y_2,y_1} - 2V_{x_1,1}V_{y_2,x_1^2} \\ d_2(V_{y_3,y_1}) &= -V_{y_2,y_1}V_{y_3,y_1} \\ d_2(V_{y_3,x_1^2}) &= -V_{x_2,x_1}V_{y_3,x_1x_2} + V_{y_2,x_1^2}V_{y_3,y_2} \\ d_2(V_{y_2,x_2}) &= V_{y_1,x_2}V_{y_2,y_1} \\ d_2(V_{y_1,x_1}) &= -V_{x_2,x_1}V_{y_1,x_2} \quad \text{and} \\ d_2(V_{x_1,1}) &= d_2(V_{y_3,y_2}) = d_2(V_{y_2,y_1}) = d_2(V_{x_2,x_1}) = d_2(V_{y_3,x_1x_2}) = d_2(V_{y_2,x_1^2}) = d_2(V_{y_1,x_2}) = 0. \end{aligned}$$

Let $V_{w_1} := 2V_{y_2,x_1} - V_{y_3,x_2}$ and $V_{w_2} := 2V_{y_2,x_2} - V_{y_1,x_1}$. Then, for $D = d_1 + d_2$, we have done from Theorem 2.1. $\square$

Lemma 5.3. *Let $M = (\Lambda V, d)$ be a minimal DGA and $m = (\Lambda U, D)$ a free DGA with $D = d_1 + d_2$. Here $d_i : U \rightarrow \Lambda^i U$. Suppose that there is a DGA-map $\varphi : M \rightarrow m$ such that $\varphi^\sharp : V \cong H^*(U, d_1)$. Then φ is a quasi-isomorphism, i.e., M is the minimal model of m .*

Proof. There are the filtrations of the word lengths on ΛV and ΛU , respectively:

$$F^i(\Lambda V) := \Lambda^{\geq i} V \quad \text{and} \quad F^i(\Lambda U) := \Lambda^{\geq i} U,$$

where $F^0 \supset \dots \supset F^n \supset F^{n+1} \supset \dots$ and $F^N H^m(\Lambda V) = F^N H^m(\Lambda U) = 0$ for a sufficient large N for each m . Note that differentials and a DGA-map preserve the filtrations. From the assumption, φ induces an isomorphism on E^1 -term of in the spectral sequence of the (decreasing) filtrations. Thus it indicates an isomorphism $\varphi^* : H^*(M) \cong H^*(m)$ from [12, Theorem 3.2] ([23, Chap.9 Theorem 3]). $\square$

Theorem 5.4. *The minimal model $M(\text{Baut}_1 X)$ of the DGA-model $m(\text{Baut}_1 X)$ is given as follows:*

$$M(\text{Baut}_1 X) \cong (\Lambda(U_{y_3,1}, U_{y_2,1}, U_{y_1,1}, U_{y_3,x_1}, U_{w_1}, U_{w_2}, U_\sigma, U_{y_1,x_2}), d)$$

where $|U_{y_3,1}| = 12$, $|U_{y_2,1}| = 10$, $|U_{y_1,1}| = 8$, $|U_{y_3,x_1}| = 8$, $|U_{w_1}| = 6$, $|U_{w_2}| = 4$, $|U_\sigma| = 3$, $|U_{y_1,x_2}| = 2$ with differential $d(U_{y_3,1}) = U_{y_2,1}U_\sigma$, $d(U_{y_2,1}) = U_{y_1,1}U_\sigma$, $d(U_{y_3,x_1}) = U_{w_1}U_\sigma$, $d(U_{w_1}) = U_{w_2}U_\sigma$, $d(U_{w_2}) = U_{y_1,x_2}U_\sigma$ and $d(U_{y_1,1}) = d(U_\sigma) = d(U_{y_1,x_2}) = 0$. Here U_σ is a non-exact cocycle corresponding to $\sigma = [(x_2, x_1) + (y_2, y_1) + 2(y_3, y_2)]$. It is not formal but coformal.

Proof. Indeed, a DGA-map $\varphi: M(\text{Baut}_1 X) \rightarrow m(\text{Baut}_1 X)$ is given as

$$\begin{aligned}
\varphi(U_{y_1, x_2}) &= V_{y_1, x_2} \\
\varphi(U_\sigma) &= V_{x_2, x_1} + 2V_{y_3, y_2} + V_{y_2, y_1} \\
\varphi(U_{w_2}) &= 2V_{w_2} + 3V_{y_2, x_1^2} V_{y_1, x_2} - 2V_{y_3, x_1 x_2} V_{y_1, x_2} \\
\varphi(U_{w_1}) &= -6V_{w_1} - 4V_{w_2} V_{y_3, x_1 x_2} + 10V_{w_2} V_{y_2, x_1^2} + 6V_{y_2, x_2} V_{y_3, x_1 x_2} - 24V_{y_2, x_2} V_{y_2, x_1^2} \\
&\quad + 6V_{y_3, x_1^2} V_{y_1, x_2} + 2V_{y_3, x_1 x_2}^2 V_{y_1 x_2} - \frac{7}{2}V_{y_2, x_1^2}^2 V_{y_1, x_2} \\
\varphi(U_{y_3, x_1}) &= -36V_{y_3, x_1} + 12V_{w_1} V_{y_3, x_1 x_2} + 6V_{w_1} V_{y_2, x_1^2} - 36V_{y_2, x_1} V_{y_3, x_1 x_2} - 72V_{y_2, x_2} V_{y_3, x_1^2} \\
&\quad + 36V_{w_2} V_{y_3, x_1^2} + 4V_{w_2} V_{y_3, x_1 x_2}^2 - 11V_{w_2} V_{y_2, x_1^2}^2 + 16V_{w_2} V_{y_3, x_1 x_2} V_{y_2, x_1^2} \\
&\quad - 12V_{y_2, x_2} V_{y_3, x_1 x_2}^2 + 24V_{y_2, x_2} V_{y_2, x_1^2}^2 - 30V_{y_2, x_2} V_{y_3, x_1 x_2} V_{y_2, x_1^2} \\
&\quad - 12V_{y_3, x_1^2} V_{y_3, x_1 x_2} V_{y_1, x_2} - 6V_{y_3, x_1^2} V_{y_1, x_2} V_{y_2, x_1^2} - \frac{4}{3}V_{y_3, x_1 x_2}^3 V_{y_1, x_2} \\
&\quad + \frac{11}{6}V_{y_2, x_1^2}^3 V_{y_1, x_2} - 6V_{y_3, x_1 x_2}^2 V_{y_1, x_2} V_{y_2, x_1^2} + V_{y_2, x_1^2}^2 V_{y_3, x_1 x_2} V_{y_1, x_2} \\
\varphi(U_{y_1, 1}) &= 3V_{y_1, 1} + 3V_{y_2, x_1} V_{y_1, x_2} + V_{w_2}^2 + 3V_{y_2, x_2}^2 - 3V_{w_2} V_{y_2, x_2} + 6V_{y_2, x_2} V_{y_1, x_2} V_{y_2, x_1^2} \\
&\quad - 2V_{w_2} V_{y_1, x_2} V_{y_2, x_1^2} + V_{y_1, x_2}^2 V_{y_2, x_1^2}^2 \\
\varphi(U_{y_2, 1}) &= 18V_{y_2, 1} - 6V_{y_1, 1} V_{y_3, x_1 x_2} + 15V_{y_1, 1} V_{y_2, x_1^2} + 9V_{y_3, x_1} V_{y_1, x_2} - 3V_{w_1} V_{w_2} + 18V_{y_2, x_1} V_{y_2, x_2} \\
&\quad - 6V_{w_1} V_{y_1, x_2} V_{y_2, x_1^2} + 3V_{y_2, x_1} V_{y_3, x_1 x_2} V_{y_1, x_2} + 15V_{y_2, x_1} V_{y_1, x_2} V_{y_2, x_1^2} - 2V_{w_2}^2 V_{y_3, x_1 x_2} \\
&\quad + 5V_{w_2}^2 V_{y_2, x_1^2} - 6V_{y_2, x_2}^2 V_{y_3, x_1 x_2} + 33V_{y_2, x_2}^2 V_{y_2, x_1^2} + 9V_{w_2} V_{y_2, x_2} V_{y_3, x_1 x_2} - 27V_{w_2} V_{y_2, x_2} V_{y_2, x_1^2} \\
&\quad - 6V_{w_2} V_{y_3, x_1^2} V_{y_1, x_2} + 18V_{y_2, x_2} V_{y_3, x_1^2} V_{y_1, x_2} - 4V_{w_2} V_{y_2, x_1^2}^2 V_{y_1, x_2} - 2V_{w_2} V_{y_3, x_1 x_2} V_{y_1, x_2} V_{y_2, x_1^2} \\
&\quad + 6V_{y_2, x_2} V_{y_2, x_1^2}^2 V_{y_1, x_2} + 12V_{y_2, x_2} V_{y_3, x_1 x_2} V_{y_1, x_2} V_{y_2, x_1^2} + 6V_{y_3, x_1^2} V_{y_1, x_2}^2 V_{y_2, x_1^2} \\
&\quad + 4V_{y_1, x_2}^2 V_{y_2, x_1^2}^2 V_{y_3, x_1 x_2} - V_{y_1, x_2}^2 V_{y_2, x_1^2}^3 \\
\varphi(U_{y_3, 1}) &= 54V_{y_3, 1} + 18V_{y_2, 1} V_{y_3, x_1 x_2} - 18V_{y_2, 1} V_{y_2, x_1^2} + 54V_{y_1, 1} V_{y_3, x_1^2} + 54V_{y_3, x_1} V_{y_2, x_2} + 18V_{w_1}^2 \\
&\quad + 54V_{y_2, x_1}^2 - 54V_{w_1} V_{y_2, x_1} + 6V_{y_1, 1} V_{y_3, x_1 x_2}^2 - \frac{33}{2}V_{y_1, 1} V_{y_2, x_1^2}^2 + 24V_{y_1, 1} V_{y_3, x_1 x_2} V_{y_2, x_1^2} \\
&\quad - 18V_{y_3, x_1} V_{y_3, x_1 x_2} V_{y_1, x_2} + 45V_{y_3, x_1} V_{y_1, x_2} V_{y_2, x_1^2} + 6V_{w_1} V_{w_2} V_{y_3, x_1 x_2} - 15V_{w_1} V_{w_2} V_{y_2, x_1^2} \\
&\quad - 36V_{w_1} V_{y_2, x_2} V_{y_3, x_1 x_2} + 36V_{w_1} V_{y_2, x_2} V_{y_2, x_1^2} + 6V_{y_2, x_2}^2 V_{y_3, x_1 x_2} V_{y_2, x_1^2} - 9V_{y_2, x_2} V_{y_2, x_1^2}^3 V_{y_1, x_2} \\
&\quad - 36V_{w_1} V_{y_3, x_1^2} V_{y_1, x_2} + 72V_{y_2, x_1} V_{y_2, x_2} V_{y_3, x_1 x_2} - 18V_{y_2, x_1} V_{y_2, x_2} V_{y_2, x_1^2} + 54V_{y_2, x_1} V_{y_3, x_1^2} V_{y_1, x_2} \\
&\quad + 108V_{y_2, x_2}^2 V_{y_3, x_1^2} - 54V_{w_2} V_{y_2, x_2} V_{y_3, x_1^2} + 6V_{w_1} V_{y_2, x_1^2}^2 V_{y_1, x_2} - 24V_{w_1} V_{y_3, x_1 x_2} V_{y_1, x_2} V_{y_2, x_1^2} \\
&\quad - 12V_{y_2, x_1} V_{y_3, x_1 x_2}^2 V_{y_1, x_2} - \frac{33}{2}V_{y_2, x_1} V_{y_2, x_1^2}^2 V_{y_1, x_2} + 69V_{y_2, x_1} V_{y_3, x_1 x_2} V_{y_1, x_2} V_{y_2, x_1^2} + 2V_{w_2}^2 V_{y_3, x_1 x_2}^2 \\
&\quad + \frac{25}{2}V_{w_2}^2 V_{y_2, x_1^2}^2 - 10V_{w_2}^2 V_{y_3, x_1 x_2} V_{y_2, x_1^2} + 24V_{y_2, x_2}^2 V_{y_3, x_1 x_2}^2 + \frac{75}{2}V_{y_2, x_2}^2 V_{y_2, x_1^2}^2 \\
&\quad + 18V_{y_3, x_1^2}^2 V_{y_1, x_2}^2 - 12V_{w_2} V_{y_2, x_2} V_{y_3, x_1 x_2}^2 - \frac{87}{2}V_{w_2} V_{y_2, x_2} V_{y_2, x_1^2}^2 + 15V_{w_2} V_{y_2, x_2} V_{y_3, x_1 x_2} V_{y_2, x_1^2} \\
&\quad + 12V_{w_2} V_{y_3, x_1^2} V_{y_3, x_1 x_2} V_{y_1, x_2} - 30V_{w_2} V_{y_3, x_1^2} V_{y_1, x_2} V_{y_2, x_1^2} + 54V_{y_2, x_2} V_{y_3, x_1^2} V_{y_1, x_2} V_{y_2, x_1^2} \\
&\quad + 5V_{w_2} V_{y_2, x_1^2}^3 V_{y_1, x_2} + 8V_{w_2} V_{y_3, x_1 x_2}^2 V_{y_1, x_2} V_{y_2, x_1^2} - 22V_{w_2} V_{y_3, x_1 x_2} V_{y_1, x_2} V_{y_2, x_1^2}^2 \\
&\quad + 36V_{y_2, x_2} V_{y_3, x_1 x_2} V_{y_1, x_2} V_{y_2, x_1^2}^2 - 6V_{y_3, x_1^2} V_{y_1, x_2}^2 V_{y_2, x_1^2}^2 + 24V_{y_3, x_1^2} V_{y_3, x_1 x_2} V_{y_1, x_2}^2 V_{y_2, x_1^2} \\
&\quad + \frac{1}{2}V_{y_1, x_2}^2 V_{y_2, x_1^2}^4 - 4V_{y_3, x_1 x_2} V_{y_1, x_2}^2 V_{y_2, x_1^2}^3 + 8V_{y_3, x_1 x_2}^2 V_{y_1, x_2}^2 V_{y_2, x_1^2}^2
\end{aligned}$$

Here we can directly check $D \circ \varphi = \varphi \circ d$. Thus we have done from Lemma 5.3. $\square$

Question 5.5. *When X is a homogeneous space, is $Baut_1X$ coformal ?*

REFERENCES

- [1] M. Amann, *Non-formal homogeneous spaces*, Math. Z. **274** (2013) 1299-1325
- [2] A. Dold and R. Lashof, *Principal quasi-fibrations and fibre homotopy equivalence of bundles*, Illinois J. Math. **3** (1959), 285-305
- [3] Y. Félix, S. Halperin and J. C. Thomas, *Rational homotopy theory*, Graduate Texts in Mathematics **205**, Springer-Verlag, 2001.
- [4] Y. Félix, G.Lupton and S.B.Smith, *Homotopy theory of function spaces and related topics*, Contemporary Math. **519** 2009.
- [5] Y. Félix and D.Tanré, *H-space structure on pointed mapping sapces*, Algebraic Geometric Topology, **5** (2005) 713-724
- [6] J.B. Gatsinzi, *The homotopy Lie algebra of classifying spaces*, J.Pure and Appl. Alg. **120** (1997) 281-289
- [7] D.H.Gottlieb, *Evaluation subgroups of homotopy groups*, Amer. J. Math., **91** (1969) 729-756
- [8] W.Greub, S.Halperin and R.Vanstone, *Connection, curvature and cohomology III*, Academic press [1976]
- [9] S.Halperin, *Finiteness in the minimal models of Sullivan*, Trans. A.M.S. **230** (1977) 173-199
- [10] S.Halperin, *Rational homotopy and torus actions*, London Math. Soc. Lecture Note Series **93**, Cambridge Univ. Press (1985) 293-306
- [11] G.Lupton and B.S.Smith, *Realizing spaces as classifying spaces*, arXiv:1502.05625v1
- [12] J. McCleary, *A user's guide to spectral sequences*, 2nd Edition, Cambridge Stud. Adv. Math., **58**, Cambridge Univ. Press, 2001.
- [13] W.Meier, *Rational universal fibrations and flag manifolds* Math. Ann. 258 (1982) 329-340
- [14] J. Milnor, *On space having the homotopy type of a CW-complex*, Trans AMS. **90** (1959) 272-280
- [15] J. Milnor, *On the characteristic classes for spherical fibrations*, Comm.Math.Helv. **43** (1968) 51-73
- [16] P.-E. Parent, *Formal and non-formal homogeneous spaces of small rank*, Univ. of Ottawa, Thesis (1996)
- [17] D. Quillen, *Rational homotopy theory*, Ann. Math. **90** (1969) 205-295
- [18] P. Salvatore, *Rational homotopy nilpotency of self-equivalences*, Topology and its Appl. **77** (1997) 37-50
- [19] M. Schlessinger and J. Stasheff, *Deformation theory and rational homotopy type*, arXiv:1211.1647v1
- [20] H. Shiga and M. Tezuka, *Rational fibrations, homogeneous spaces with positive Euler characteristics and Jacobians*, Ann. Inst. Fourier (Grenoble) **37** (1987) 81-106
- [21] S.B. Smith, *Rational type of classifying spaces for fibrations*, Contemporary Math. **274** (2001) 299-307
- [22] S.B. Smith, *The rational homotopy Lie algebra of classifying spaces for formal two-stage spaces*, J.P.A.A. **160** (2001) 333-343
- [23] E.H. Spanier, *Algebraic Topology*, Springer G.T.M. 1966
- [24] D.Sullivan, *Infinitesimal computations in topology*, I.H.E.S., **47** (1978) 269-331
- [25] D. Tannré, *Homotopie Rationnelle: Modèles de Chen, Quillen, Sullivan*, Lecture Note in Math. Springer **1025** (1983)
- [26] T. Yamaguchi, *When is the classifying space for elliptic fibrations rank one ?* Bull. Korean Math **42** (2005) 521-525

KOCHI UNIVERSITY, 2-5-1, KOCHI,780-8520, JAPAN
E-mail address: cosmo51mutta@yahoo.co.jp

KOCHI UNIVERSITY, 2-5-1, KOCHI,780-8520, JAPAN
E-mail address: tyamag@kochi-u.ac.jp